

Referral Status:	MRN:
<input type="checkbox"/> New referral	<input type="checkbox"/> Order change
<input type="checkbox"/> Order Renewal	
Patient preferred clinic:	

Skyrizi® (risankizumab-rzaa) Standard Plan of Treatment for Ulcerative Colitis

PATIENT DEMOGRAPHICS:

Date of Referral:	Patient's Phone:
Patient Name:	Address:
Date of Birth:	City, State, Zip:
Height in inches:	Weight: LB or KG
Gender:	Allergies:
	See list
	NKDA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

K51.0 - Ulcerative (Chronic) Pancolitis	K51.4 - Inflammatory Polyps of Colon
K51.2 - Ulcerative (Chronic) Proctitis	K51.5 - Left Sided Colitis
K51.3 - Ulcerative (Chronic) Rectosigmoiditis	K51.8 - Other Ulcerative Colitis or Unspecified
- Other:	

REQUESTED DOCUMENTATION:

PREVIOUS ADMINISTRATION: HAS THIS PATIENT TAKEN THIS MEDICATION BEFORE?

1	Insurance information	IF NO:	IF YES:
2	Most recent History & Physical	PLEASE STATE	LAST INFUSION DATE:
3	Full medication list	REQUIRED WASHOUT	NEXT INFUSION DATE:
4	Tried and failed therapies	FROM PREVIOUS	IF ORDER CHANGE:
5	REQUIRED: TB screening for new start	THERAPY:	
6	Baseline LFTs and bilirubin level		
			Continue current order until insurance approved

MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive risankizumab-rzaa if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new-onset or deterioration neurological changes, and/or surgery.

PREMEDICATION TO BE ADMINISTERED 30 MINUTES PRIOR TO ADMINISTRATION AS SELECTED

IV	Diphenhydramine	25mg	50mg	Other:	PO	Acetaminophen	325mg	500mg	650mg	1000mg
	Methylprednisolone	40mg	125mg			Famotidine	20mg	40mg		
	Famotidine	20mg	40 mg			Diphenhydramine	25mg	50mg		
	Other:					Fexofenadine	60mg	180mg		
						Cetirizine	10mg			
				Loratadine	10mg					
				Other:						

MEDICATION/DOSE:

Skyrizi® (risankizumab-rzaa) 1200mg in 250ml or 500ml of NS given IV over at least 2 hours

SPECIAL/OTHER LAB ORDERS:

FREQUENCY:

Week 0, week 4, and week 8

Other: _____

LINE USE/CARE ORDERS:

- Start PIV/Access CVC
- Flush device per facility standard flushing procedure

ADVERSE REACTION & ANAPHYLAXIS ORDERS:

Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.



PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures)

DATE:

Dispense as written/Brand medically necessary	Substitution permitted