Щ	Palmetto
NON	TNELISTON®

INFUSION® Phone: 1-800-809-1265 Fax: 1-866-872-8920

Referral Status:					
	New referral				
Patient preferred clinic:					

MRN: Order change

Order Renewal

	iximab Unspec		an of Tr	eatment for D)erm	natology					
PATIENT DEMOGRAPHICS:											
Date of Referral:				Patient's Phone:							
Patient Name:				Address:							
Date of Birth:					City, State, Zip:						
	it in inches:	Weight:			Gender: Allergies: See list NKDA						
DIAC	GNOSIS: (PLEASE CO	OMPLETE	2 ^{mb} AND		MPLE	TE ICD 10 FOR BI	ILLING)				
	L40.5 Psoriatic	Arthritis/Arth	ropathy								
	L40 Psoriasis	;									
	Other:										
REQ	UESTED DOCUMEN	ITATION:				ATION: HAS THIS	PATIENT TA	AKEN THIS	MEDICATIO	N	
1	Insurance information			IF NO:	IF YE						
2	Most recent History & F	Physical		PLEASE STATE REQUIRED WASHOUT	LAST INFUSION DATE:						
3	Full medication list			FROM PREVIOUS							
4	Tried and failed therapi			THERAPY:	IF ORDER CHANGE:						
5	REQUIRED: TB screen	0	•	-	Continue current order until insurance approved						
6	REQUIRED: HBsAg, a		anti-HBs								
	ICATION ORDERS:		6					and infection			
	Patient may be ineligible to rec w-onset or deterioration neur		-		process,	, antifungal therapy, active f	ever and/or suspe	ected infection, r	iew or worsening	symptoms of	
	EDICATION TO BE AL		-		NISTR	RATION AS SELECTE	ED				
Prem	edication with antihistam	nines, acetan	ninophen, ai	nd/or corticosteroids m	nay be	considered to preven	t infusion-relat	ed reactions.			
	Diphenhydramine	25mg	50mg			Acetaminophen	325mg	500mg	650mg	1000mg	
IV	Methylprednisolone	40mg	125mg	Other:		Famotidine	20mg	40mg			
	Famotidine	20mg	40 mg			Diphenhydramine	25mg	50mg			
	Other:				PO	Fexofenadine	60mg	180mg			
SPEC	CIFIC MEDICATION:					Cetirizine	10mg				
Remicade Any infliximab						Loratadine	10mg				
	Avsola					Other:					
					FRE	REQUENCY:					
Renflevis <u>useu accorui</u>					Induction to be completed at week 0, week 2, and week 6, and						
DOSE:				nes	then every 8 weeks thereafter						
	5mg/kg diluted in NS infused IV per step protocol			ol over 2 hours	Maintenance every 8 weeks						
	Other:					Infuse every weeks					
	May utilize expedited in	nfusion per p	protocol to r	un over 1 hour as tole	rated						
*1611.	1					CIAL/LAB ORDER	ς.				
	kimab doses <1000mg)mg in 1000ml NS (max			-		7	<u>.</u>				
~2000		x concentrat	uon–4mg/n	II)							
Pres	criber confirms that the	patient has	been evalua	ted and screened for t	the pre	esence of hepatitis B	virus (HBV) pr	ior to initiatir	na treatment.	 Prescriber to	
				ptoms of HBV and TB							
						Refills x 12 months unless noted otherwise here:					
LINE	USE/CARE ORDER					ADVERSE REACTION & ANAPHYLAXIS ORDERS:					
\checkmark	Start PIV/Access CV					Administer acute infusion and anaphylaxis					
Flush device per facility standard flushing procedure						medications per Palmetto Infusion standing					
Provide nursing care per Palmetto Infusion Nursing Procedure					es	adverse reaction orders, which can be found at our website or scan here.					
	and post procedure		n if indicate	d						Children III I	
	SCRIBER INFORMA	TION:									
PROVIDER NAME:					PHONE:						
ADDRESS:						FAX:					
CITY, STATE, ZIP:						NPI:					
PRESCRIBER SIGNATURE: (No stamp signatures)								DATE:			
	Dispense as wr	itten/Brand	medically	necessary			Substitution	n permitted			