Palmetto INFUSION®

Phone: 1-800-809-1265 Fax: 1-866-872-8920

Referral Status:		MRN:
☐ New referral	☐ Order change	☐ Order Renewal
Patient preferred clinic:		

Infliximab Unspecified Plan of Treatment for Gastroenterology

PATIENT DEMOGRAPHICS:

Date of Referral:		Patient's Phone:	
Patient Name:		Address:	
Date of Birth:		City, State, Zip:	
Height in inches:	Weight: LB or KG	Gender:	Allergies: ☐ See list ☐ NKDA

DIAGNOSIS: (PLEASE COMPLETE 2ND AND 3RD DIGITS TO COMPLETE ICD 10 FOR BILLING)

☐	K50.0___ - Crohn's Disease (small intestine)	☐	K51.5___ - Left sided Ulcerative (chronic) Colitis
☐	K50.1___ - Crohn's Disease (large intestine)	☐	K51.0___ - Universal Ulcerative (chronic) Pancolitis
☐	K50.8___ - Crohn's Disease (small & large intestine)	☐	K51.8___ - Other Ulcerative (chronic) Colitis
☐	K50.9___ - Crohn's Disease, unspecified	☐	K60.3___ - Anal Fistula
☐	K63.2- Fistula of Intestine	☐	_____ - Other:

REQUESTED DOCUMENTATION:

1	Insurance information
2	Most recent History & Physical
3	Full medication list
4	Tried and failed therapies
5	**REQUIRED:** TB screening for new start
6	**REQUIRED**: HBsAg, anti-HBc, and anti-HBs

PREVIOUS ADMINISTRATION: HAS THIS PATIENT TAKEN THIS MEDICATION BEFORE?

IF NO: PLEASE STATE REQUIRED WASHOUT FROM PREVIOUS THERAPY:

IF YES:
LAST INFUSION DATE:
NEXT INFUSION DATE:

IF ORDER CHANGE:

☐ **Continue current order until insurance approved**

MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive infliximab if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new or worsening symptoms of CHF, new onset or deterioration neurological changes, and/or surgery.

PREMEDICATION TO BE ADMINISTERED 30 MINUTES PRIOR TO ADMINISTRATION AS SELECTED

Premedication with antihistamines, acetaminophen, and/or corticosteroids may be considered to prevent infusion-related reactions.

IV

Diphenhydramine	☐ 25mg	☐ 50mg		
Methylprednisolone	☐ 40mg	☐ 125mg	☐ Other:	
Famotidine	☐ 20mg	☐ 40 mg		
Other:	☐			

PO

Acetaminophen	☐ 325mg	☐ 500mg	☐ 650mg	☐ 1000mg
Famotidine	☐ 20mg	☐ 40mg		
Diphenhydramine	☐ 25mg	☐ 50mg		
Fexofenadine	☐ 60mg	☐ 180mg		
Cetirizine	☐ 10mg			
Loratadine	☐ 10mg			
Other:	☐			

SPECIFIC MEDICATION:

- ☐ Remicade
- ☐ Avsola
- ☐ Inflectra
- ☐ Renflexis

☐ **Any infliximab biosimilar may be used according to payer guidelines**

DOSE/RATE:

- ☐ 5mg/kg diluted in NS infused IV per step protocol over 2 hours
- ☐ Other:_____
- ☐ Expedited infusion rate of 1 hour if tolerated

Infliximab doses <1000mg in 250ml NS, doses >1000mg in 500ml NS and doses >2000mg in 1000ml NS (max concentration=4mg/ml)

FREQUENCY:

- ☐ Induction to be completed at week 0, week 2, week 6, and then every 8 weeks thereafter
- ☐ Maintenance every 8 weeks
- ☐ Infuse every _____ weeks

SPECIAL/LAB ORDERS:

☐ _____

_____

Prescriber confirms that the patient has been evaluated and screened for the presence of hepatitis B virus (HBV) prior to initiating treatment. Prescriber to monitor patient for symptoms of HBV and TB infection and reactivation as clinically appropriate.

☑ Refills x 12 months unless noted otherwise here:

NURSING ORDERS:

- ☑ Start PIV/Access CVC
- ☑ Flush device per facility standard flushing procedure
- ☑ Provide nursing care per Palmetto Infusion Nursing Procedures and post procedure observation if indicated

ADVERSE REACTION & ANAPHYLAXIS ORDERS:

☑ Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.

PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures)		DATE:
Dispense as written/Brand medically necessary	Substitution permitted	