

Referral Status:	MRN:
<input type="checkbox"/> New referral	<input type="checkbox"/> Order change
<input type="checkbox"/> Order Renewal	
Patient preferred clinic:	

## Simponi ARIA® (golimumab) Standard Plan of Treatment for Rheumatology

### PATIENT DEMOGRAPHICS:

Date of Referral:	Patient's Phone:						
Patient Name:	Address:						
Date of Birth:	City, State, Zip:						
Height in inches:	Weight:	LB or	KG	Gender:	Allergies:	See list	NKDA

### DIAGNOSIS: (PLEASE COMPLETE 2<sup>ND</sup> AND 3<sup>RD</sup> DIGITS TO COMPLETE ICD 10 FOR BILLING)

M05._____ - Rheumatoid Arthritis with Rheumatoid factor	M06._____ - Rheumatoid Arthritis without Rheumatoid factor
L40.5_____ - Psoriatic Arthropathy	M45._____ - Ankylosing Spondylitis
_____ - Other:	

### REQUESTED DOCUMENTATION: PREVIOUS ADMINISTRATION: HAS THIS PATIENT TAKEN THIS MEDICATION BEFORE?

1	Insurance information	IF NO:	IF YES:
2	Most recent History & Physical	PLEASE STATE	LAST INFUSION DATE:
3	Full medication list	REQUIRED WASHOUT	NEXT INFUSION DATE:
4	Tried and failed therapies	FROM PREVIOUS	<b>IF ORDER CHANGE:</b> <div style="border: 1px solid black; padding: 5px; text-align: center;"> <b>Continue current order until insurance approved</b> </div>
5	<b>REQUIRED:</b> TB screening for new start	THERAPY:	
6	HBV screening/labs as required by payor		

### MEDICATION ORDERS:

NOTE: Patient may be ineligible to receive golimumab if receiving antibiotics for active infectious process, antifungal therapy, active fever and/or suspected infection, new or worsening symptoms of CHF, new onset or deterioration neurological changes, and/or surgery

#### PREMEDICATION TO BE ADMINISTERED 30 MINUTES PRIOR TO ADMINISTRATION AS SELECTED

<b>IV</b>	Diphenhydramine	25mg	50mg	Other:	<b>PO</b>	Acetaminophen	325mg	500mg	650mg	1000mg
	Methylprednisolone	40mg	125mg			Famotidine	20mg	40mg		
	Famotidine	20mg	40 mg			Diphenhydramine	25mg	50mg		
	Other:					Fexofenadine	60mg	180mg		
						Cetirizine	10mg			
				Loratadine	10mg					
				Other:						

**MEDICATION/DOSE:**

Simponi ARIA® (golimumab) 2 mg/kg per 100 ml NS given IV to infuse over at least 30 minutes

**FREQUENCY:**

Induction: Given at 0 week and 4 weeks, and then every 8 weeks thereafter

Maintenance: Given every 8 weeks

Other: \_\_\_\_\_

**SPECIAL/OTHER LAB ORDERS:**


\_\_\_\_\_

\_\_\_\_\_

Prescriber confirms that the patient has been evaluated and screened for the presence of hepatitis B virus (HBV) prior to initiating treatment. Prescriber to monitor patient for symptoms of HBV and TB infection and reactivation as clinically appropriate.

Refills x 12 months unless noted otherwise here:

### LINE USE/CARE ORDERS: ADVERSE REACTION & ANAPHYLAXIS ORDERS:

<input checked="" type="checkbox"/> Start PIV/Access CVC <input checked="" type="checkbox"/> Flush device per facility standard flushing procedure	Administer acute infusion and anaphylaxis medications per Palmetto Infusion standing adverse reaction orders, which can be found at our website or scan here.	
---	---	---

### PRESCRIBER INFORMATION:

PROVIDER NAME:	PHONE:
ADDRESS:	FAX:
CITY, STATE, ZIP:	NPI:

PRESCRIBER SIGNATURE: (No stamp signatures)	DATE:
Dispense as written/Brand medically necessary	Substitution permitted